



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

## NEW LITERATURE.

BY W. A. KELLERMAN.

"UEBER LEPTOSPHERIA NIGRANS" (Rob. et Derm.) L. Fuckelii Niessl und verwante Arten. J. Niessl. Hedwigia, Mar. u. Apr., 1887.

"NACHTRÄGE UND BERICHTIGUNGEN ZU SACCARDO'S SYLLOGE FUNGORUM, VOL. I, II." Von Dr. G. Winter.

"SOME AUSTRALIAN FUNGI." By M. C. Cooke. Grevillea, June, 1887.

"NEW BRITISH FUNGI." By M. C. Cooke l. c.

"NEW SPECIES OF RAVENELIA." l. c.

"RAVENELIA VERRUCOSA, Cke. & Ellis—Hypophylla; uredosporis, globosis, asperulis, luteis (16  $\mu$ ); teleutosporis in glomerulos hæmisphæricos congestis (80  $\mu$ ); glomerulis (sporis 20), stipitatis, cum lobulis (circa 8), hyalinos circumdatis; telutosporis cuneatis ad apicem asperulis, atrofusis (20  $\mu$  diam.) On leaves of *Lecania* sp.? Mexico (J. B. Ellis.)

It differs from *R. stictica* in not being sessile, in the hyaline lobules being larger and more conspicuous and the warts smaller. This is the only species with which it could be confounded, and from this it seems to be distinct.

"SOME NEW BRITISH DISCOMYCETES." By W. Phillips, F. L. S. l. c.

"BRITISH PYRENUMYCETES. A PRELIMINARY LIST OF KNOWN SPECIES." By G. Masee; continued. l. c.

"SYNOPSIS PYRENUMYCETUM." Continued. l. c.

"HEMIARCYRIA CHRYSOSPORA, LISTER." By Arthur Lister. l. c.

"FUNGUS OF ANEMONE BEDS. PEZIZA TUBEROSA, BULL." Worthington G. Smith, Gardener's Chronicle, May 28, 1887.

## CORRECTIONS.

In North American Fungi, Cent. XIX (No. 1889), *Ustilago* "*subinclusa*," Kuhn, should be *U. subinclusa*, Kornicke.

No. 1869 is no doubt what Fckl. in his Enumeratio-Fungorum Nassoriæ published as *Uromyces Prunorum*, but, as Dr. Winter has pointed out to us, is not a true *Uromyces* but only the uredo form of *Puccinia Prunorum*, Lk., to which Fuckel himself (Symb. Myc., p. 49) afterwards referred it.

No. 1878. At the time this was issued, we overlooked the fact that it had already been distributed under another name, *Chrysomyxa albida*, Kuhn. See Rab. Winter Fung. Euro., No. 3015.

The following corrections to be made in this JOURNAL: On p. 52, 7th line from top, for *Erigerontes* read *Erigerontis*; on p. 53, 7th line from top, for *paniflora* read *pauciflora*; on p. 53, 13th line from top, for *Cuphua* read *Cuphea*; on p. 57, 2d line from bottom, for *Cerryana* read *Therryana*; on p. 69, 8th line from bottom, for *Phododendri* read *Rhododendri*.